

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application.

1. (Previously presented) A method for enhancing the efficacy of a therapeutic treatment for cancer in a patient, said therapeutic treatment being selected from the group consisting of: chemotherapy, radiation therapy, surgery, and combinations thereof, said method comprising the steps of:

administering to said patient a therapeutically effective amount of a carbohydrate which binds to a galectin; and

administering said therapeutic treatment to said patient.

2. (Original) The method of claim 1, wherein said galectin is present on the cell surface of a tissue of said patient.

3. (Previously presented) The method of claim 1, wherein said carbohydrate binds to galectin-1 or galectin-3.

4. (Previously presented) The method of claim 1, wherein said carbohydrate comprises a polymeric backbone having side chains dependent therefrom, said side chains being terminated by a galactose or arabinose unit.

5. (Previously presented) The method of claim 1, wherein said carbohydrate comprises a substantially demethoxylated polygalacturonic acid which is interrupted with rhamnose residues.

6. (Cancelled)

7. (Previously presented) The method of claim 1, wherein said carbohydrate comprises a branched carbohydrate.

8. (Previously presented) The method of claim 1, wherein said carbohydrate comprises a modified pectin.

9. (Original) The method of claim 8, wherein said modified pectin comprises a pH modified pectin.

10. (Original) The method of claim 9, wherein said modified pectin comprises an enzymatically modified pectin.

11. (Original) The method of claim 8, wherein said modified pectin comprises a thermally modified pectin.

12. (Original) The method of claim 8, wherein said modified pectin comprises a modified citrus pectin.

13. (Previously presented) The method of claim 1, wherein said carbohydrate has a molecular weight of at least 300 dalton.

14. (Previously presented) The method of claim 1, wherein said carbohydrate has a molecular weight in the range of 300-2,000 dalton.

15. (Original) The method of claim 8, wherein said modified pectin has a molecular weight in the range of 1-50 kilodalton.

16. (Original) The method of claim 8, wherein said modified pectin has a molecular weight in the range of 1-15 kilodalton.

17. (Original) The method of claim 8, wherein said modified pectin has a molecular weight of approximately 10 kilodalton.

18. (Previously presented) The method of claim 1, wherein said step of administering said carbohydrate to said patient comprises injecting said carbohydrate into said patient.

19. (Previously presented) The method of claim 1, wherein said step of administering said carbohydrate to said patient comprises orally administering said carbohydrate to said patient.

20. (Previously presented) The method of claim 1, wherein said step of administering said carbohydrate to said patient comprises administering said carbohydrate prior to administering said therapeutic treatment to said patient.

21. (Previously presented) The method of claim 1, wherein said step of administering said carbohydrate to said patient comprises administering said carbohydrate to said patient after said therapeutic treatment is administered to said patient.

22. (Previously presented) The method of claim 1, wherein said carbohydrate is administered concomitant with said therapeutic treatment.

23. (Previously presented) A method for enhancing the efficacy of a therapeutic treatment for cancer in a patient, said therapeutic treatment being selected from chemotherapy, radiation therapy, surgery, and combinations thereof, said method comprising
administering to said patient a therapeutically effective amount of a carbohydrate which binds to a galectin; and
administering said therapeutic treatment to said patient,
wherein said carbohydrate comprises polymeric backbone having side chains dependent therefrom.

24. (Previously presented) The method of claim 23, wherein said polymeric backbone comprises homopolymer.

25. (Previously presented) The method of claim 23, wherein said carbohydrate binds to galectin-3.

26. (Previously presented) The method of claim 23, wherein said carbohydrate is a naturally occurring carbohydrate or a modified product thereof.

27. (Previously presented) A method for enhancing the efficacy of a surgical treatment for cancer in a patient, said method comprising

administering to said patient a therapeutically effective amount of a carbohydrate comprising a polymeric backbone having side chains dependent therefrom, said side chains being terminated by a galactose or arbinose unit, and administering surgery to said patient.

28. (Previously presented) A method for enhancing the efficacy an oncolytic chemotherapeutic in a patient, said method comprising

administering to said patient, prior to or concomitant with said oncolytic chemotherapeutic, a therapeutically effective amount of a carbohydrate comprising a polymeric backbone having side chains dependent therefrom, said side chains being terminated by a galactose or arabinose unit, and administering said oncolytic chemotherapeutic to said patient.